

Mark J. Coronel, MD

496 County Road 111 • Building D • Suite 2 • Manorville, NY 11949
34 Commerce Drive • Riverhead, NY 11901
Tel: 631-591-3000 • Fax: 631-591-1734
www.eastendgastro.com

POST-OP INSTRUCTIONS

Restriction On Activity

- Do not drive, return to work or operate machinery until the day after the procedure.
- Following day return to usual activity, unless otherwise instructed.
- No lifting, straining or running for 24 hours.
- Diet avoid alcohol for 24 hours after procedure; otherwise eat and drink as usual.
- Do not make any important, personal or business decisions, sign any legal papers, or perform and activity that depends on your full concentrating power or mental judgment.
- Do not drink any alcohol or take any nerve or sleeping drugs as they may add to the effects of medicine still present in your body.

Treatment For Common After Effects

- Mild abdominal pain and bloating Rest and take liquids only.
- Sore throat- treat with throat lozenges or gargle with salt water.

Symptoms To Watch For And Report To Your Physician

- Chills or fever occurring within 24 hours after procedure.
- Persistent rectal bleeding.
- Severe continuing abdominal pain, bloating, vomiting or nausea.
- Persistent IV site swelling / redness.
- Severe pain in chest or neck
- IV site stays red and swollen for more than 2 days
- Persistent black, tarry bowel movements.

If A Polyp Has Been Removed, For The Next 14 Days

- Consult your physician before taking any medications.
- If rectal bleeding occurs, call your physician.