

PLEASE READ YOUR PACKET



EAST END GASTROENTEROLOGY & HEPATOLOGY

Blood Thinners: Hold 48 hours Prior to procedure.

COLO

COLO/ENDO

w/MOVIprep

PATIENT: _____ **DATE SCHEDULED:** _____

APPOINTMENT TIME: _____ **ARRIVAL TIME:** _____

Location: 287 Wading River Road Unit 2, Manorville, NY 11949

Directions to our Manorville Location:

GPS / GOOGLE MAPS – LI Urgent Care

From LIE Heading East: Take the LIE to Exit 69, south on Wading River Road, Continue South over South Street. Head south towards Sunrise Highway. Manorville Square is located on the left- hand side of Wading River Road before Sunrise highway, across from the Animal Farm Petting Zoo. We share the same parking lot as LI Urgent Care.
From LIE Heading West: Take LIE to Exit 69, Make a Left onto Wading River Road, Continue South over South Street heading towards Sunrise highway, Manorville square is located on the left-hand side of Wading River Road across from the Animal Farm Petting zoo.

From Sunrise Highway Heading West: Take Sunrise Highway to Exit 59, we are located in the same parking lot as LI Urgent care right off of Sunrise highway North service road located behind the EXXON gas station. **From Sunrise Highway Heading EAST:** Take Sunrise Highway Exit 59 for Wading River Road/ County RD 25 towards Center Moriches- Make a Left at Wading River Road Light(7 Eleven will be to your right) Cross over Sunrise Hwy Bridge, EXXON Gas Station will be on the right. Manorville Square entrance will be on the right-hand side after EXXON.

PLEASE NOTE:

**PATIENTS THAT CANCEL WITH LESS THAN 72 HOURS
NOTICE WILL INCUR A \$100 CANCELLATION FEE**

PLEASE READ YOUR PACKET

COLONOSCOPY INSTRUCTIONS w/ MOVIPREP

You have been scheduled for a colonoscopy. This information sheet will inform you of how to best prepare for the procedure. If there are any questions remaining after reading these instructions, please call the office.

If you are NOT able to keep your appointment, please give us at least 72 hours notice.

IMPORTANT NOTES:

1. **You must arrange to have someone accompany you for the procedure.** You will NOT Be allowed to drive home, and you may NOT drive for the rest of the day. On the day of the Procedure, if you arrive without someone to take you home, your procedure will NOT be done.
2. You cannot go to work the day of the procedure.
3. If you take any BLOOD THINNERS such as Coumadin, Plavix, Effient, Warfrin or Pradaxa on a daily basis, you must notify the Doctor immediately. These medications may need to be discontinued PRIOR to your colonoscopy.
4. BLOOD PRESSURE Medications may be taken VERY early with a small sip of water.
5. You must thoroughly read the instructions on the following pages.

COLONOSCOPY INFORMATION:

Colonoscopy is a diagnostic procedure that uses a thin flexible tube with a camera and a light to examine the lining of your large intestine (known as the colon). During the exam, you will be lying on your left side on a stretcher. An intravenous catheter (an "IV") will be placed in a vein in your arm. You will then be hooked up to monitoring equipment to measure your heart rate, blood pressure and oxygen level of your blood. An Anesthesiologist or his/her assistant will administer medicine into your I.V. for sedation (***please notify us of any allergies to medicines***). Once you are asleep and comfortable, the colonoscope will be gently inserted into the rectum and carefully advanced through the colon. If a colonic polyp, an abnormal growth of colonic tissue is found, your physician may be able to remove the polyp through the instrument. In rare cases, passage of the colonoscope through the entire colon cannot be achieved. The entire examination usually lasts from 20 minutes to one hour.

The **benefits** of the procedure include the ability to detect and remove colon polyps for colon cancer prevention, evaluate gastrointestinal and abdominal symptoms, obtain biopsies, and identify and treat sites of bleeding. Although Colonoscopy is a safe procedure, complications can occur. The **Risks** include but are not limited to, Bleeding which can occur after polyp removal or therapy and generally stops on

PLEASE READ YOUR PACKET

its own but on occasion may require blood transfusion, repeat colonoscopy, hospitalization and/or surgery ; Perforation which is a puncture or tear in colon wall which may require hospitalization, surgery and may be serious complication that can even cause death; Reaction to sedation which very rarely some patients develop low blood pressure, irregular heartbeat or difficulty breathing; Missed polyps or lesions which, because the test is not perfect, can occur particularly if they are small in size or flat in shape.

After the procedure, you might be sleepy for one to two hours after the procedure due to medications you will receive at the time of the procedure. You may expel some gas as air, which is instilled into the colon during the colonoscopy, for adequate visualization. You may experience some abdominal bloating and distention for several hours after the colonoscopy. You will be able to resume your diet after the colonoscopy, unless you are instructed otherwise.

ONE WEEK PRIOR TO EXAM:

- **Arrange for someone to drive you home from your procedure. YOU WILL NOT BE ALLOWED TO DRIVE HOME OR TAKE A TAXI WITHOUT AN ADULT (OTHER THAN THE TAXI DRIVER) TO ACCOMPANY YOU. On the day of the procedure, if you Arrive without someone to take you home, your procedure WILL NOT be done.**
- **Arrange to take the day of the procedure off from work. You cannot go to work; you are to stay at home the remainder of the day after the procedure and relax. You may not DRIVE FOR THE REST OF THE DAY!**
- **Stop any vitamins containing iron and/or iron supplements one week prior to the Colonoscopy.**
- **Purchase your prep!**

THREE DAYS PRIOR

- **Please limit intake of corn, celery, lettuce, nuts and seeds as these are poorly digestible and often leave residue in the colon and can interfere with exam visualization.**

DAY BEFORE PROCEDURE

- **START CLEAR LIQUID DIET** for the ENTIRE day before the procedure which means
- **NO SOLID FOOD, NO GUM, NO MILK OR MILK PRODUCTS, NO JUICE WITH PULP OR SEEDS.**
- **YOU MAY ONLY HAVE THE FOLLOWING:**
Water, Seltzer, Apple Juice, White Grape Juice, Sprite, Ginger Ale, 7 UP, Mountain Dew, light colored sports drinks, crystal light, green or white tea, Bouillon or Clear Broth, Coffee, Tea, NO MILK, Gatorade (NO RED), Italian Ice (NO RED), Plain Jello (NO RED), Popsicles (NO DAIRY or pops with fruit pieces).
There are NO exceptions, patients may eat and drink ONLY WHAT IS ON THIS LIST!!
- **BEGIN YOUR PREP AS DIRECTED.**

PLEASE READ YOUR PACKET

MOVIPREP INSTRUCTIONS

Pick up from the pharmacy

1. Moviprep prescription which includes 4 pouches and one disposable container for mixing.
2. 1 64 oz. or 2 32oz. bottles of orange Gatorade

THE DAY BEFORE THE PROCEDURE:

1. **CLEAR LIQUID DIET ALL DAY. NO SOLID FOOD.**
2. **BEFORE 4PM (usually in morning)-** Empty one pouch A and one pouch B of the MoviPrep into the disposable container. Add orange Gatorade to the top fill line of the Container. Mix to dissolve. Refrigerate!
3. **AT 6PM-** Start drinking the solution. The MoviPrep container is divided into 4 marks. Drink Down to each line (approximately 8 oz) every 15 minutes (6:00, 6:15, 6:30, 6:45, 7:00). Afterwards, drink at least 2 additional glasses of any clear liquid as described above.
3. **AFTER 7 PM (ONCE FINISHED DRINKING FROM CONTAINER)-** Empty one pouch A and one pouch B of the MoviPrep into the disposable container. Add orange Gatorade to the top fill line of the Container. Mix to dissolve. Refrigerate!

THE DAY OF THE PROCEDURE:

1. **You should take your blood pressure and heart medications early in the morning with a small sip of water.** If you are diabetic, please see diabetic instruction sheet or call the office for any questions.
2. **4-5 hours before you leave your house for the procedure drink the other container.** Take the MoviPrep container from the refrigerator, drink the solution about 8 oz. every 15 minutes until you have completed the full container, then drink 16 oz. of the clear liquid of your choice. **THIS MUST BE COMPLETED AT LEAST 4 HOURS PRIOR TO YOUR COLONOSCOPY.** If your procedure is scheduled early in the morning, you will need to get up in the night to take this dose of preparation. **Do not take anything by mouth for 4 hours prior to your colonoscopy.**
3. **HAVE NOTHING ELSE TO EAT OR DRINK, NOT EVEN WATER, until after your procedure.**

Due to the anticipated large volume of diarrhea, you can use baby wipes for cleaning and the use of Destin or Vaseline for skin protection.

DO NOT OMIT ANY PART OF THE PREPARATION. A POOR PREPARATION MAY CAUSE A

PLEASE READ YOUR PACKET

POLYP OR CANCER TO GO UNDETECTED.